JEFFERS S N Invasive Species



Scentless chamomile

(Tripleurosperum inodorum)



Scentless chamomile				
General	Family: Sunflower (Asteraceae) Introduced from Europe Colorado List B – Containment required			
Common names	Mayweed, scentless mayweed			
Habitat	Forb Can live as an annual, biennial or short-lived perennial Found in meadows, pastures, rights-of-way, drainages			
PLANT				
Vegetation	Plant to 3 feet tall Leaves ferny, odorless			
Roots	Fibrous			
Flower	June-October White with yellow center Centers tend to dome and the petals bend down as the flowers mature Single on each stem 3/4 - 1 inch wide			
Seed	As many as 300,000 per plant May last up to 15 years in the soil			
Seedling	Spring May form dense patches			
Reproduction	Seed			

CONTROL				
BIOLOGICAL	None			
CHEMICAL	Stage	Herbicide	Timing	
	Rosette to bolting	Aminopyralid (Milestone) Chlorsulfuron (Telar) Metsulfuron (Escort XP)	Spring- early Summer	
CULTURAL	Prevention Removal - make sure the entire root is removed. Must be repeated as new plants germinate Shallow tillage when plants are small, before flowers form			
MECHANICAL	Stage	Method	Timing	
Burning	N/A			
Grazing	N/A	May cause blistering to livestock		
Mowing	N/A	Plants will regrow		

Use all chemicals according to the manufacturer's label.

No specific recommendation or endorsement is made or implied by listing the above methods or products.

1/2021